TOOL: Walk-in Medical Assistant Blood Pressure Check Protocol (Kaiser Permanente)

KAISER FOUNDATION HOSPITAL SOUTHERN CALIFORNIA PERMANENTE MEDICAL GROUP POLICY AND PROCEDURE

Section		NUMBER	
OPE	RATIONS	NUMBER	2008
Title	Title	EFFECTIVE: DATE	9/91
BLOOD PRESSURE MONITORING - Blood Pressure Check Visit	REVISION	11/09, 1/11	
		PAGE NUMBER	1 of 4

PURPOSE

To promote health maintenance and improve patient care through appropriate blood pressure monitoring.

POLICY

- All adult patients presenting for an appointment or procedure will have their blood pressure measured when instructed to do so by the Proactive Office Encounter automated alert
- 2. Take and document a second blood pressure when indicated
- 3. Give the After Visit Summary (AVS) to all patients

AUTHORIZED PERSONNEL

- 1. Provider MD, PA, NP, CNM, CRNA
- 2. Pharm D
- 3. RN
- 4. LVN/MA/Ortho Tech

EQUIPMENT/SUPPLIES NEEDED

- 1. Sphygmomanometer or Automated Blood Pressure Machine
- 2. Appropriate-sized blood pressure cuff
- Stethoscope
- 4. Chair with back support
- 5. Table to support arm at heart level

PROCEDURE

*Refer to Vital Signs P&P

Adult Primary Care Departments (scheduled provider visit):

- Using appropriate technique, take the patient's sitting BP and enter in KPHealthConnect™ in the "Vitals Section". The date and time will automatically populate. After each documented measurement and at completion of vitals documentation click the "Close" button in the "Vitals" section to ensure capture of all values.
- The HTN "Best Practice Alert" will populate when the BP is elevated: greater than 139 systolic and/or greater than 89 diastolic instructing the staff to repeat the blood pressure.
- 3. If the BP is elevated, wait one (1) minute and repeat. If the patient is 70 years of age or older take the second blood pressure in the standing position. Enter the new BP in KPHealthConnect™ in the "New Set of Vital Signs" section. The date and time will automatically populate. Click the "Close" button in the "Vitals" section. If the BP was standing, click on "Doc Flowsheet" in upper right corner. Under the recent BP, in the "BP Patient Position" box, select the paper icon, select standing then "Accept".
- 4. Inform the member of their blood pressure.
- 5. The HTN "Best Practice Alert" will be addressed by the provider.

Module Blood Pressure Check scheduled appointment or walk-in (provider visit not scheduled):

Note: As with any visit, if the patient expresses any complaints or questions, the MA must refer the patient to the RN or Provider

- 1. Open the patient's chart from the resources schedule using Allied Health Encounter.
- Using appropriate technique, take the patient's sitting BP and enter in KPHealthConnect™ in the "Vitals Section". The date and time will automatically populate. After each documented measurement and at completion of vitals documentation click the "Close" button in the Vitals section to ensure capture of all values.
- The HTN "Best Practice Alert" will populate when the BP is elevated: greater than 139 systolic and/or greater than 89 diastolic instructing the staff to repeat the blood pressure.

TOOL: Walk-in Medical Assistant Blood Pressure Check Protocol (Kaiser Permanente)

KAISER FOUNDATION HOSPITAL SOUTHERN CALIFORNIA PERMANENTE MEDICAL GROUP POLICY AND PROCEDURE

Section	1	AUUMDED	
	OPERATIONS	NUMBER	2008
Title	BLOOD PRESSURE MONITORING - Blood Pressure Check Visit	EFFECTIVE: DATE	9/91
		REVISION	11/09, 1/11
		PAGE NUMBER	2 of 4

- 4. If the BP is elevated, wait one (1) minute and repeat. If the patient is 70 years of age or older take the second blood pressure in the standing position. Enter the new BP in KPHealthConnect™ in the "New Set of Vital Signs" section. The date and time will automatically populate. Click the "Close" button in the "Vitals" section. If the BP was standing, click on "Doc Flowsheet" in upper right corner. Under the recent BP, in the "BP Patient Position" box, select the paper icon, select standing then "Accept".
- 5. Inform the member of their blood pressure.
- 6. Follow procedure in Table A based on the lowest BP reading.

Specialty Care Departments:

- Using appropriate technique, take the patient's sitting BP and enter in KPHealthConnect™ under the "Vitals Section". Click the "Close" button in the Vitals section.
- 2. The HTN "Best Practice Alert" will populate when the BP is elevated: greater than 139 systolic and /or greater than 89 diastolic instructing the staff to repeat the blood pressure.
- 3. If the BP is elevated, wait one (1) minute and repeat. If the patient is 70 years of age or older take the second blood pressure in the standing position. Enter the new BP in HealthConnect™ in the "New Set of Vital Signs" Section. Click the "Close" button in the Vitals section. If the BP was standing, click on "Doc Flowsheet" in upper right corner. Under the recent BP, in the "BP Patient Position" box, select the paper icon, select standing then "Accept".
- 4. If BP greater than 139/89, open the nursing note to document the elevated BP.
- 5. Inform the member of their Blood Pressure
- 6. Forward encounter to PCP by opening Follow up
 - 7.1. Add PCP to recipient list
 - 7.2. Enter .HBP in Routing Comments
 - 7.3. Exit the work space
- If NO PCP assigned (this will not be consistent in every medical center) refer to HTN Hotline, if one is available, or PC DOD.

ADULT BLOOD PRESSURE CATEGORIES and ACTIONS (Table A)

CATEGORY	SBP	DBP	Action
Low∙	109 or less		 MA/LVN: Recheck BP with patient standing. Standing SBP is 110 or greater, follow Action in "Controlled" category in this table. Standing SBP is 109 or less, refer to RN for evaluation. Release patient as directed by RN or provider with BP recheck appointment in 2 months. Close encounter. RN: SBP 100 – 109 without symptoms and previous SBP 110 or greater: Instruct staff to make follow up BP check appointment in 2 months, route encounter to PCP with smart phrase ".vs" in "Routing Instructions" and release pt. SBP 100 – 109 without symptoms and previous SBP 100 – 109 without symptoms and previous SBP 100 – 109, discuss with provider for possible medication adjustment and timeframe for follow up BP check.

TOOL: Walk-in Medical Assistant Blood Pressure Check Protocol (Kaiser Permanente)

KAISER FOUNDATION HOSPITAL SOUTHERN CALIFORNIA PERMANENTE MEDICAL GROUP POLICY AND PROCEDURE

Section OPERATIONS	NUMBER	2008
BLOOD PRESSURE MONITORING - Blood Pressure Check Visit	EFFECTIVE: DATE	9/91
	REVISION	11/09, 1/11
	PAGE NUMBER	3 of 4

			 SBP 100 – 109 with symptoms or SBP 99 or less without symptoms, discuss with provider for possible medication adjustment and timeframe for follow up BP check. SBP 99 or less with symptoms, instruct staff to schedule pt to see provider today
Controlled ·	110-139	≤ 89	MA/LVN: Direct the patient to follow-up with another blood pressure check in about 6 months. Release the patient. Close encounter
Stage 1 Hypertension	140-159	90-99	MA/LVN: Depending on module workflow Make BP recheck appointment in 2-4 weeks and release the patient. Route to PCP using ".HBP" in "Routing Instructions" OR Inform provider pt is waiting for further instructions OR Inform module RN that pt is waiting for further instructions
Stage 2 Hypertension	160 179	100 - 109	MA/LVN: Refer patient to RN or provider for evaluation prior to releasing the patient. Release the patient as directed by the RN or provider. Schedule the patient for a BP recheck in 1-2 weeks. Route to PCP using ".HBP" in "Routing Instructions" RN: No symptoms: Instruct MA/LVN to release patient with instructions above. Symptoms: Instruct MA to schedule pt to see provider today
Stage 2 Hypertension Urgent	<u>></u> 180	<u>></u> 110	Refer to provider for evaluation. Do not release the patient.

SmartPhrase List: Suggested RN Documentation Phrases (Table B)

Stage 1 hypertension

.BP4 - BP 140-159/90-99

Lowest BP 140-159/90-99. Blood pressure elevated. Current treatment to be continued. The staff is instructed to please inform patient of his/her blood pressure today. Tell patient to continue current mediations and that today's visit will be routed to their PCP for review. We will notify them if doctor orders a change in medication. Prior to patient leaving, schedule appointment for blood pressure recheck in 2-4 weeks. The PCP was messaged with his/her last 3 encounter BP readings.

Stage 2 hypertension: RN to discuss case with provider to determine if medication adjustment needed prior to release of patient

.BP2 - BP 160-179/100-109

Lowest BP 160-179/100-109. The staff is instructed to schedule an appointment for blood pressure recheck in 1-2 weeks before he/she leaves. Please inform patient of his/her blood pressure today. Tell patient to continue

TOOL: Walk-in Medical Assistant Blood Pressure Check Protocol (Kaiser Permanente)

KAISER FOUNDATION HOSPITAL SOUTHERN CALIFORNIA PERMANENTE MEDICAL GROUP POLICY AND PROCEDURE

Section	OPERATIONS	NUMBER	2008
Title	BLOOD PRESSURE MONITORING - Blood Pressure Check Visit	EFFECTIVE DATE	9/91
		REVISION	11/09, 1/11
		PAGE NUMBER	4 of 4

current medications and that today's visit will be routed to their PCP for review. PCP was messaged with his/her last 3 encounter BP readings...

.BPTODAY - BP ≥180/110

Lowest Systolic BP greater than or equal to 180 and/or Diastolic BP greater than or equal to 110. His/Her blood pressure is elevated, requiring evaluation today. The staff is instructed to schedule him/her to see a provider for blood pressure evaluation today.

MARE

Please have patient come in for an MA blood pressure check in *** week(s).

Low Blood Pressure

.BPlowOK - SBP 100 - 109 standing asymptomatic on antihypertension medications

Systolic BP 100 – 109 standing, he/she is asymptomatic and he/she is on hypertension medications. His/Her PCP was messaged with his/her last 3 encounter BP readings. The staff is instructed to schedule him/her an appointment with his/her PCP for follow-up in three months.

.BPlow – 1) Systolic BP less than 100 standing, symptomatic or asymptomatic or (2) Systolic BP 100 – 109 standing and symptomatic The staff is instructed to schedule him/her to see a provider for blood pressure evaluation today.